



¿Se emplea con rigor científico la terminología ecológica en las publicaciones sobre la transmisión trófica de patógenos?

GONZÁLVEZ, Moisés; MOLEÓN, Marcos; MARTÍNEZ-CARRASCO, Carlos

INTRODUCCIÓN

MATERIAL Y MÉTODOS


RESULTADOS

DISCUSIÓN

CONCLUSIONES

Ciencia basada en...


Ciencia basada en...



```
graph TD; A([Ciencia basada en...]) --> B[Revisión bibliográfica contrastadas];
```

Revisión bibliográfica
contrastadas


Ciencia basada en...





```
graph TD; A([Ciencia basada en...]) --> B[Revisión bibliográfica contrastadas]; A --> C[Evidencias elaboradas con rigor científico];
```


Evidencias elaboradas
con rigor científico

Revisión bibliográfica
contrastadas









Pullin *et al.*, 2004



Rigor científico

Transmisión de patógenos por consumo de carne en fauna silvestre


CONSUMO DE CARROÑA
DEPREDACIÓN

Intraespecífica


Interespecífica



Terminología ecológica




Colaboraciones entre disciplinas



Van Noorden, 2015

Colaboraciones entre disciplinas



Objetivos

En relación a las publicaciones científicas sobre transmisión de patógenos por consumo de carne en fauna silvestre...

¿Cuál es la disciplina a la que pertenecen los autores?

¿Se emplea correctamente la terminología ecológica en estos trabajos?

INTRODUCCIÓN

MATERIAL Y MÉTODOS

RESULTADOS

DISCUSIÓN

CONCLUSIONES

Base consultada: WoS

Web of Science™
InCites™
Journal Citation Reports®
Essential Science Indicators™
EndNote™
Iniciar sesión
Ayuda
Español

WEB OF SCIENCE™
THOMSON REUTERS™

Buscar
Colección principal de Web of Science™
Mis herramientas
Historial de búsqueda
Lista de registros marcados

¡Bienvenido al nuevo Web of Science! [Vea un breve tutorial.](#)

Búsqueda básica

Ejemplo: oil spill* mediterranean
Tema
Buscar

[+ Agregar otro campo](#) | [Borrar todos los campos](#)

Haga clic aquí para obtener sugerencias para mejorar su búsqueda.

PERÍODO DE TIEMPO

☐ Todos los años
☒ Desde 1900 hasta 2016

▶ MÁS AJUSTES

Periodo de búsqueda: Junio-Julio 2016

Base consultada: WoS

Web of Science™ InCites™ Journal Citation Reports™

WEB OF SCIENCE™

Buscar Colección principal de Web of Science

Búsqueda básica

Ejemplo: oil spill* mediterranean


PERÍODO DE TIEMPO

☐ Todos los años

☒ Desde 1900 hasta 2016

MÁS AJUSTES

Sin restricción temporal



- ☐ INFECTIOUS DISEASES (6,397)
- ☒ PARASITOLOGY (5,583)
- ☐ ZOOLOGY (4,891)
- ☒ VETERINARY SCIENCES (3,811)
- ☐ IMMUNOLOGY (3,545)
- ☐ BIOCHEMISTRY MOLECULAR BIOLOGY (2,764)
- ☐ PHYSIOLOGY (2,677)
- ☐ ENVIRONMENTAL SCIENCES ECOLOGY (1,895)
- ☒ PATHOLOGY (1,665)
- ☐ GASTROENTEROLOGY HEPATOLOGY (1,581)
- ☐ DEVELOPMENTAL BIOLOGY (1,330)
- ☐ PUBLIC ENVIRONMENTAL OCCUPATIONAL HEALTH (1,302)
- ☐ PHARMACOLOGY PHARMACY (1,290)
- ☐ ANATOMY MORPHOLOGY (1,255)
- ☐ GENETICS HEREDITY (1,110)
- ☐ CELL BIOLOGY (1,020)
- ☐ FOOD SCIENCE TECHNOLOGY (987)
- ☐ CARDIOVASCULAR SYSTEM CARDIOLOGY (938)

Búsqueda básica

Ejemplo: oil spill* mediterranean

PERÍODO DE TIEMPO

☒ Todos los años

☐ Desde 1900 hasta 2017

MÁS AJUSTES


Sugerir de forma automática nombres

☒ Activada



Haga clic aquí para obtener sugerencias para mejorar su búsqueda.

Áreas seleccionadas: “PARASITOLOGY”, VETERINARY SCIENCE” y “PATHOLOGY”




“Carrion”

“Meat”

“Scaveng*”

“Predat*”

+ “Pathogen” OR “Trichinella” OR “Toxoplasma” OR “Neospora”



“Carrion”

“Meat”

“Scaveng*”

“Predat*”

+ “Pathogen” OR “Trichinella” OR “Toxoplasma” OR “Neospora”

Web of Science™ InCites™ Journal Citation Reports® Essential Science Indicators™ EndNote™

WEB OF SCIENCE™ THOMSON REUTERS™

Mis herramientas ▾ Historial de búsqueda Lista de registros marcados

Búsqueda básica ▾

Ejemplo: oil spill* medline

Agrege otro campo | Borrar todos los campos

Buscar

PERÍODO DE TIEMPO

Desde 1900 hasta 2017

MÁS AJUSTES

Sugerir de forma automática nombres de publicaciones

Activada ▾

“Carrion”

“Meat”

+ “Pathogen” OR “Trichinella” OR “Toxoplasma” OR “Neospora”

“Scaveng*”

“Predat*”

Web of Science™ InCites™ Journal Citation Reports® Essential Science Indicators™ EndNote™

WEB OF SCIENCE™ THOMSON REUTERS™

Mis herramientas ▾ Historial de búsqueda Lista de registros marcados

Búsqueda básica ▾

Ejemplo: oil spill* medline

Buscar

PERÍODO DE TIEMPO

Desde 1900 hasta

MÁS AJUSTES

Sugerir de forma automática nombres de publicaciones

Activada ▾

“Carrion”

“Meat”

+ “Pathogen” OR “Trichinella” OR “Toxoplasma” OR “Neospora”

“Scaveng*”

“Predat*”

Web of Science™ InCites™ Journal Citation Reports® Essential Science Indicators™ EndNote™

WEB OF SCIENCE™ THOMSON REUTERS™

Mis herramientas ▾ Historial de búsqueda Lista de registros marcados

Búsqueda básica ▾

Ejemplo: oil spill* medline

Buscar

PERÍODO DE TIEMPO

Desde 1900 hasta

MÁS AJUSTES

Sugerir de forma automática nombres de publicaciones

Activada ▾


“Carrion”

“Meat”

+ “Pathogen” OR “Trichinella” OR “Toxoplasma” OR “Neospora”

“Scaveng*”

“Predat*”




12919
artículos


**12919
artículos**


**2603
artículos**

Filtro áreas: "Veterinary Science",
"Pathology" y "Parasitology"

10316







Transmisión de patógenos por consumo de carne



Review

ELSEVIER

Journal homepage: www.elsevier.com/locate/jcp

Review

Implications of wild dog ecology on the sylvatic and domestic life cycle of *Neospora caninum* in Australia

Y. King^{a,e}, David J. Jenkins^b, John T. Ellis^c, Peter Fleming^d, Peter A. Windsor^a, Ja...

^a - Sydney, NSW 2006, Australia
^b - Sydney, NSW 2006, Australia
^c - Wagga Wagga, NSW 2678, Australia
^d - Wagga Wagga, NSW 2677, Australia
^e - Sydney, NSW 2000, Australia

of Neoplasia

Jessica S. King^{a,e}, David J. Jenkins^a, ...
Wagga Wagga, NSW 2678, Australia
Wagga Wagga, NSW 2607, Australia
Wagga Wagga, NSW 2680, Australia
Wagga Wagga, NSW 2600, Australia
Wagga Wagga, NSW 2601, Australia

^a Faculty of Veterinary Science, University of Sydney, NSW 2006, Australia
^b School of Animal and Veterinary Science, Charles Sturt University, Wagga Wagga, NSW 2607, Australia
^c Department of Medical and Molecular Biosciences, University of Technology, Sydney, NSW 2000, Australia
^d Department of Medical and Molecular Biosciences, University of Technology, Sydney, NSW 2000, Australia
^e Veterinary Pest Research Unit, Industry & Investment NSW, Orange Agricultural Institute, ACT 2601, Australia
^f Invasive Animals Cooperative Research Centre, University of Canberra, Canberra, ACT 2601, Australia

ABSTRACT

... either trans

ARTICLE INFO

Article history:
Accepted 2 March 2010

Keywords:
Neosporosis
Horizontal transmission

ABSTRACT

ABSTRACT

Neospora caninum is transmitted either transplacentally or horizontally in tissues or oocysts shed by dogs. Neosporosis is a significant problem in cattle, as infected cows may remain infective for up to 7 months of pregnancy. Infected cows may remain infective for 1 or more consecutive or non-consecutive pregnancies. A great deal is known about the disease, although only limited information is available on the maintenance of the parasite in Australia, the presence of the dingo as the top-order predator.

ScienceDirect



Toxoplasma gondii prevalence in Israeli crows and Griffon


H. Salant^{a,*}, J. Hamburger^a, R. King^b, G. Ganeth^c

ARTICLE INFO

Article history:
Received 17 June 2012
Received in revised form 12 July 2012
Accepted 27 July 2012

Keywords:
Toxoplasma gondii
MAT
Crows
Griffon vultures
Israel

ABSTRACT

[illegible]

Red fox (*Vulpes vulpes*) cannibalistic behaviour and the prevalence of *Trichinella britovi* in NW Italian Alps

Received: 24 June 2005 / Accepted: 28 July 2005 / Published online: 7 September 2005
© Springer-Verlag 2005


Introduction

Abstract Food habits of the red fox (*Vulpes vulpes*) were studied in the Aosta Valley region (NW Italian Alps) and were related to the prevalence of Trichinellosis infection in the red fox itself and in two mammalian species (the stone marten (*Martes foina*) and the European badger (*Meles meles*). The search of Trichinella by automatic digestion of muscle samples and the determination of the prevalence of Trichinella by usages of 7.9 ± 4.3% in stone martens and 1.9 ± 1.8% in red foxes, with significant differences between the two species. All larvae were identified among the species *T. britovici*. This diet was identified as *Trichinella britovici*. The killed animals ($n=180$) and the stomach contents analysis of road-killed animals ($n=109$). Our results confirmed an opportunistic feeding behaviour of the red fox, which is able to use various trophic resources, both animal and vegetal origin: e.g. wild and cultivated fruit ($\% = 47.1$; $\% = 67.3$), roe deer ($\% = 64.8$) and brown bear ($\% = 15.6$; $\% = 22.8$) formed the bulk of the fox's diet. The frequency of occurrence of Trichinella in the fox's diet was lower than in the stone marten. The frequency of

In the temperate regions of Eurasia, three *Trichinella* parasitic sylvatic cycle, *T. britovi*, found mainly in wild boar and horse meat, and *T. spiralis*, found mainly in wild boar and in mammals and birds (Pozio 2001), even if the systematic of the genus is still discussed (Bessonov *et al.* 2002; Murrel *et al.* 2000).

The sylvatic cycle is supposed to involve mainly carnivores with cannibalistic and scavenger behaviour, the eating of carrion being considered the principal mechanism of nematode transmission (Campbell 1988). In Italy, researches focused on the red fox (*Vulpes vulpes*), which is actually considered the main or almost unique reservoir of the sylvatic cycle of the parasite (Pozio 1991; Rossi *et al.* 1992; Rossi and Balbo 1993).

While there are various researches about the distribution and prevalence of the nematode in different wild animal species (see, for example, Campbell 1988 and Pozio




Molecular epidemiology of *Trichinella* spp. in three Baltic countries: Lithuania, Latvia, and Estonia

A. Malakauskas • V. Paulauskas • T. Järvis • P. Keidans •
C. Eddi • C. M. O. Kapel

Received: 12 May 2006 / Accepted: 24 August 2006 / Published online: 30 September 2006
© Springer-Verlag 2006

Abstract Meat of domestic pigs is the significant source of emerged human trichinellosis in Lithuania, Latvia, and Estonia over the past two decades. However, there is very little known on the occurrence of *Trichinella* spp. in main wildlife reservoirs and its transmission in domestic and sylvatic cycles in these countries. The present study demonstrated considerably higher endemicity of *Trichinella* spp. in main sylvatic reservoirs (28.9–42% in foxes (*Vulpes vulpes*) and raccoon dogs (*Nyctereutes procyonoides*) in all three countries than

previously reported. Molecular analysis of the larvae from more than 500 samples of meat revealed four *Trichinella* species: *Trichinella britovi*, *Trichinella pseudospiralis* (synonym *Trichinella spiralis*) several as the first record of nonencapsulating *T. pseudospiralis* in the Eastern European region (domestic pigs in Lithuania and Poland respectively) and only *Trichinella spiralis* was detected in the



The role of wild carnivores in the maintenance of trichinellosis in the Tatras National Park, Slovakia

Z. HURNÍKOVÁ¹, B. CHOVANCOVÁ², D. BARTKOVÁ³, P. DUBINSKÝ¹

¹Parasitological Institute of Slovak Academy of Sciences, Hlinkova 3, 040 01 Košice, Slovak Republic, E-mail: hurnik@saske.sk; ²Research Centre of the Tatras National Park, 059 60 Tatranská Lomnica, Slovak Republic

Summary

A survey on trichinellosis in wild carnivores and their role in the transmission of parasite in the model territory of the Tatras National Park was conducted in years 2005–2006. Muscle samples of 102 wild predators representing 10 species belonging to families Canidae, Mustelidae and Ursidae were examined. The highest prevalence of *Trichinella britovi* was established among red foxes (19.7%) and martens (37.5%). The parasite was also detected in European sylvatic animals.

increase of red fox population in recent years provides a potential for spread of *Trichinella* spp. in wildlife. Whereas recent data on trichinellosis in other carnivore species are lacking, we aimed to study the presence of parasite in those animal species, inhabiting the territory of protected areas of the Tatras National Park (TANAP).

Materials and Methods

Análisis artículos

Transmisión de patógenos por consumo de carne



Available online at www.sciencedirect.com

ScienceDirect

Veterinary Parasitology 153 (2008) 152–156

veterinary
parasitology

www.elsevier.com/locate/vetpar

Short communication

Prevalence of antibodies against *Toxoplasma gondii* in roe deer from Spain

J.A. Gamarra^a, O. Cabezón^b, M. Pabón^b, M.C. Arnal^c, D.F. Luco^c,
J.P. Dubey^{d,*}, C. Gortázar^a, S. Almeria^{b,c}

^a Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13071, Ciudad Real

^b Parasitology and Parasitic Diseases, Veterinary School, Autonomous University of Barcelona, Barcelona, Spain

^c SEDIFAS, Veterinary School, University of Zaragoza, Zaragoza, Spain

^d Animal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD 20705, USA

^{*} Research Centre in Animal Health (CRESA), Autonomous University of Barcelona, Barcelona, Spain

Received 10 August 2007; received in revised form 8 January 2008; accepted 11 January 2008

Valoración individual de cada artículo

Clasificación disciplina de autores

Clasificación empleo de terminología

CONSUMO DE CARNE

DEPREDACIÓN

CARROÑO

DEPREDACIÓN
INTRAESPECÍFICA
O CANIBALISMO

DEPREDACIÓN
INTERESPECÍFICA

CARROÑO
INTRAESPECÍFICO
O CANIBALISMO

CARROÑO
INTERESPECÍFICO

Análisis artículos

Transmisión de patógenos por consumo de carne



Available online at www.sciencedirect.com

ScienceDirect

Veterinary Parasitology 153 (2008) 152–156

veterinary
parasitology

www.elsevier.com/locate/vetpar

Short communication

Prevalence of antibodies against *Toxoplasma gondii* in roe deer from Spain

J.A. Gamarra^a, O. Cabezón^b, M. Pabón^b, M.C. Arnal^c, D.F. Luco^c,
J.P. Dubey^{d,*}, C. Gortázar^a, S. Almeria^{b,c}

^a Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13071, Ciudad Real

^b Parasitology and Parasitic Diseases, Veterinary School, Autonomous University of Barcelona, Barcelona, Spain

^c SEDIFAS, Veterinary School, University of Zaragoza, Zaragoza, Spain

^d Animal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD 20705, USA

^{*} Research Centre in Animal Health (CRESA), Autonomous University of Barcelona, Barcelona, Spain

Received 10 August 2007; received in revised form 8 January 2008; accepted 11 January 2008

Valoración individual de cada artículo

Clasificación disciplina de autores

Clasificación empleo de terminología

CONSUMO DE CARNE

DEPREDACIÓN

CARROÑEO

DEPREDACIÓN
INTRAESPECÍFICA
O CANIBALISMO

DEPREDACIÓN
INTERESPECÍFICA

CARROÑEO
INTRAESPECÍFICO
O CANIBALISMO

CARROÑEO
INTERESPECÍFICO

Análisis artículos

Transmisión de patógenos por consumo de carne



Available online at www.sciencedirect.com



Veterinary Parasitology 153 (2008) 152–156

veterinary
parasitology

www.elsevier.com/locate/vetpar

Short communication

Prevalence of antibodies against *Toxoplasma gondii* in roe deer from Spain

J.A. Gamarra^a, O. Cabezón^b, M. Pabón^b, M.C. Arnal^c, D.F. Luco^c,
J.P. Dubey^{d,*}, C. Gortázar^a, S. Almeria^{b,c}

^a Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13071, Ciudad Real

^b Parasitology and Parasitic Diseases, Veterinary School, Autonomous University of Barcelona, Barcelona, Spain

^c SEDIFAS, Veterinary School, University of Zaragoza, Zaragoza, Spain

^d Animal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD 20705, USA

* Research Centre in Animal Health (CRESA), Autonomous University of Barcelona, Barcelona, Spain

Received 10 August 2007; received in revised form 8 January 2008; accepted 11 January 2008

Valoración individual de cada artículo

Clasificación disciplina de autores

Clasificación empleo de terminología

CONSUMO DE CARNE

DEPREDACIÓN

CARROÑEO


DEPREDACIÓN
INTRAESPECÍFICA
O CANIBALISMO

DEPREDACIÓN
INTERESPECÍFICA

CARROÑEO
INTRAESPECÍFICO
O CANIBALISMO

CARROÑEO
INTERESPECÍFICO

Clasificación de los artículos (disciplina de autores)



Análisis de los términos científicos empleados



The Veterinary Journal 188 (2011) 24–33

Contents lists available at ScienceDirect

The Veterinary Journal

journal homepage: www.elsevier.com/locate/tvj



ns of wild dog ecology on the sylvatic and domestic life cycle a caninum in Australia

g^{a,e}, David J. Jenkins^b, John T. Ellis^c, Peter Fleming^d, Peter A. Windsor^a, Jan Šlapeta^{a,*}

^aScience, University of Sydney, NSW 2006, Australia
^bVeterinary Science, Charles Sturt University, Wagga Wagga, NSW 2678, Australia
^cand Molecular Biosciences, University of Technology, Sydney, NSW 2007, Australia
^dArch Unit, Industry & Investment NSW, Orange Agricultural Institute, NSW 2800, Australia
^eoperative Research Centre, University of Canberra, Canberra, ACT 2601, Australia

ABSTRACT

Neospora caninum is transmitted either transplacentally or horizontally by ingestion of tissue cysts present in tissues or oocysts shed by dogs. Neosporosis is a significant disease, causing cattle abortion at 5–7 months of pregnancy. Infected cows may remain infective for life transmitting the infection in several consecutive or non-consecutive pregnancies. A great deal is known about the epidemiology of neosporosis, although only limited information is available on the main routes of horizontal transmission. In Australia, the presence of the dingo as the top-order predator suggests a potential sylvatic route of



Available online at www.sciencedirect.com

ScienceDirect

Veterinary Parasitology 153 (2008) 152–156

veterinary
parasitology

www.elsevier.com/locate/vetpar

Short communication

Prevalence of antibodies against *Toxoplasma gondii* in roe deer from Spain

J.A. Gamarra^a, O. Cabezón^b, M. Pabón^b, M.C. Arnal^c, D.F. Luco^c,
J.P. Dubey^{d,*}, C. Gortázar^a, S. Almería^{b,c}

^aInstituto de Investigación en Recursos Cinegéticos (ISREC-UCLM-JCCM), Ronda de Toledo s/n, 13071, Ciudad Real, Spain

^bParasitology and Parasitic Diseases, Veterinary School, Autonomous University of Barcelona, Barcelona, Spain

^cSEDIAS, Veterinary School, University of Zaragoza, Zaragoza, Spain

^dAnimal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Agricultural Research Service,

United States Department of Agriculture, Beltsville, MD 20705, USA

^{*}Research Centre in Animal Health (CRESA), Autonomous University of Barcelona, Barcelona, Spain

Received 10 August 2007; received in revised form 8 January 2008; accepted 11 January 2008

Abstract

Roe deer (*Capreolus capreolus*) is an important game animal in Spain. Sera from 278 roe deer from eight areas in mainland Spain were assayed for antibodies to *Toxoplasma gondii* by modified agglutination test (MAT). Titers of 1:25 or higher were found in 109 (39.2%) of 278 deer. No significant differences in antibody prevalence were found between sex or age categories. In contrast, significant differences in seroprevalence between locations were evident. Roe deer from the Northern coastal habitats (high humidity and roe deer density) had the highest prevalence, compared with low prevalence in Central Spain (arid areas and low roe deer density). There was a positive correlation between antibody prevalence and mean annual rainfall ($r_s = 0.85$, $n = 8$, $P < 0.01$). These findings have environmental and/or public health implications because venison can be an important meat source of *T. gondii* infections for humans and food animals.

© 2006 Parasitological Institute of SAS, Košice
DOI 10.2478/s11687-006-0050-9

HELMINTHOLOGIA, 44, 1: 18–20, JANUARY 2007

The role of wild carnivores in the maintenance of trichinellosis in the National Park, Slovakia

Z. HURNÍKOVÁ¹, B. CHOVANCOVÁ², D. BARTKOVÁ¹, P. DUBINSKY¹

¹Parasitological Institute of Slovak Academy of Sciences, Hlinkova 3, 040 01 Košice, Slovak Republic
²Research Centre of the Tatras National Park, 059 60 Tatranská Lomnica, Slovakia

Summary

A survey on trichinellosis in wild carnivores and their role in the transmission of parasite in the model territory of the Tatras National Park was conducted in years 2005–2006. Muscle samples of 102 wild predators representing 10 species belonging to families Canidae, Mustelidae and Ursidae were examined. The highest prevalence of *Trichinella britovi* was established among red foxes (19.7%) and martens (37.5%). The parasite was also detected in Euro-

increase of red fox population in recent years. The potential for spread of *Trichinella* spp. in these areas recent data on trichinellosis in other carnivore species are lacking, we aimed to study the presence of these animal species, inhabiting the territory of the Tatras National Park (TANAP).

Materials and Methods




Mapas bibliométricos que representan:



Mapas bibliométricos que representan:

- **Relación entre términos científicos mediante clusters.**





Mapas bibliométricos que representan:

- Relación entre términos científicos mediante clusters.

- **Evolución temporal en el empleo de estos términos.**







Mapas bibliométricos que representan:


- Relación entre términos científicos mediante clusters.
- Evolución temporal en el empleo de la terminología.
- **Relación de citas entre todos los conceptos.**

WEB OF SCIENCE™


Clasificación de los artículos (terminología)

Material y Métodos





¿Cuál es el grado de interrelación entre términos ecológicos y epidemiológicos?



INTRODUCCIÓN

MATERIAL Y MÉTODOS

RESULTADOS

DISCUSIÓN

CONCLUSIONES

Transmisión de patógenos por consumo de carne

Disciplinas de los autores en los artículos analizados

	Veterinarios	Biólogos	Médicos	Otros
Veterinarios	68			
	2			
Biólogos	14	1		
Médicos	4	1	1	
Otros	1	1	0	0

Transmisión de patógenos por consumo de carne

Disciplinas de los autores en los artículos analizados

	Veterinarios	Biólogos	Médicos	Otros
Veterinarios	68			
	2			
Biólogos	14	1		
Médicos	4	1	1	
Otros	1	1	0	0

Transmisión de patógenos por consumo de carne

Disciplinas de los autores en los artículos analizados

	Veterinarios	Biólogos	Médicos	Otros
Veterinarios	68			
	2			
Biólogos	14	1		
Médicos	4	1	1	
Otros	1	1	0	0

Transmisión de patógenos por consumo de carne

Disciplinas de los autores en los artículos analizados

	Veterinarios	Biólogos	Médicos	Otros
Veterinarios	68			
	2			
Biólogos	14	1		
Médicos	4	1	1	
Otros	1	1	0	0

Transmisión de patógenos por consumo de carne

Disciplinas de los autores en los artículos analizados

	Veterinarios	Biólogos	Médicos	Otros
Veterinarios	68			
	2			
Biólogos	14	1		
Médicos	4	1	1	
Otros	1	1	0	0

Evaluación disciplinas autores

Resultados

Unidisciplinarios

75,3%

Multidisciplinarios


24,7%

Unidisciplinarios

75,3%


Multidisciplinarios

24,7%




Unidisciplinarios

75,3%




Multidisciplinarios

24,7%




Transmisión de patógenos por consumo de carne

Evolución disciplinas/años




Transmisión de patógenos por consumo de carne

Evolución disciplinas/años



Transmisión de patógenos por consumo de carne


Evolución disciplinas/años



Transmisión de patógenos por consumo de carne

Empleo de la terminología en artículos

■ Correcto (impreciso) ■ No emplea ■ Correcto (preciso) ■ Erróneo




Unidisciplinares (75,3%)

Multidisciplinares (24,7%)


Unidisciplinar


■ Correcto (impreciso) ■ No emplea ■ Correcto (preciso) ■ Erróneo




Multidisciplinar


■ Correcto (impreciso) ■ No emplea









RELACIÓN ENTRE TÉRMINOS ECOLÓGICOS Y EPIDEMIOLÓGICOS (MAPA DE CALOR)







INTRODUCCIÓN

MATERIAL Y MÉTODOS

RESULTADOS


DISCUSIÓN

CONCLUSIONES

Falta colaboración entre equipos multidisciplinares


Falta colaboración entre equipos multidisciplinares

Multitud factores ecológicos y epidemiológicos



Falta colaboración entre equipos multidisciplinares


Multitud factores ecológicos y epidemiológicos




Falta conocimiento en áreas de estudio complejas

Falta colaboración entre equipos multidisciplinares

Multitud factores ecológicos y epidemiológicos




Falta conocimiento en áreas de estudio complejas




Terminología incorrecta/imprecisa



¿Cuáles son las ventajas de la colaboración interdisciplinar?



¿Cuáles son las ventajas de la colaboración interdisciplinar?



¿Cuáles son las ventajas de la colaboración interdisciplinar?



INTRODUCCIÓN

MATERIAL Y MÉTODOS

RESULTADOS

DISCUSIÓN

CONCLUSIONES

- 1. Escasa colaboración multidisciplinar en estudios de transmisión de patógenos por consumo de carne en fauna silvestre.**
- 2. En las publicaciones elaboradas por equipos unidisciplinares no se suelen usar términos ecológicos, y cuando se emplean es habitual que sean imprecisos o incluso erróneos.**
- 3. Para mejorar el impacto de las publicaciones epidemiológicas en este campo, es necesario que exista una mayor colaboración multidisciplinar.**



Muchas gracias

Grazie mille

Merci beaucoup